

TITLE: Moran Th-U TN250

AUTHOR(S):

S. B. Moran
Roger P. Kelly
Graduate School of Oceanography,
University of Rhode Island
Narragansett, RI, 02882
Tel : (401) 874-6530
Fax: (401) 874-6811
Email: moran@gso.uri.edu, rokelly@gso.uri.edu

FUNDING SOURCE AND GRANT NUMBER

NSF: ARC-0732680

DATA SET OVERVIEW:

The data in the accompanying file (Moran Th-U TN250.xls) includes data corresponding to samples collected by the Moran Geochemistry Lab on the spring leg of the BEST-BSIERP study from 18 June 2010 to 12 July 2010 aboard the R/V Thomas G. Thompson. Data are organized by station and sample depth, with station position (latitude and longitude) also included.

This dataset includes the following: Total ^{234}Th , and dissolved ^{238}U activities. These radionuclides are reported in units of dpm (disintegrations per minute) L^{-1} . In addition are the deployment and recovery dates and positions.

INSTRUMENT DESCRIPTION:

Samples were collected using 30-L Niskin bottles from the Thompson's CTD-rosette.

DATA COLLECTION and PROCESSING:

Total ^{234}Th was measured on water samples collected from the Thompson's CTD-rosette. Samples were typically collected from the 'hydro' cast, and bottles were sampled after gasses and nutrients were collected. Water samples were drained into graduated 4L bottles. 8-10 drops of ~30% $\text{NH}_4(\text{OH})$ was then added to raise the pH to ~9. Then, 25 μl of 0.2M KMnO_4 and 11 μl of 1 M MnCl_2 were added to form a MnO_2 suspension. This suspension stood for one hour to scavenge dissolved ^{234}Th , and was subsequently filtered onto a 1 μm (nominal) GMF filter. The filters were then dried at 60°C overnight, and then mounted for beta counting. These samples were analyzed for ^{234}Th using a RISØ GM-25-5 beta counter at sea for the initial counts, and at URI-GSO for the subsequent

counts (up to 180 days after collection). Consult Buessler et al 1992 and Buessler et al 2001 for further details.

Dissolved ^{238}U was calculated using a known relationship with salinity:

$$^{238}\text{U} (\text{dpm L}^{-1}) = S \times 0.069$$

This relationship was established in Chen et al 1986. Salinity values from the CTD's primary conductivity sensor recorded in the bottle data files were used for this calculation.

DATA FORMAT:

Data are presented as an excel file named Moran Th-U TN250.xls, as it includes the total thorium and dissolved uranium data from the BEST summer cruise of the Thompson in 2010.

Data are organized by cruise ID, station number, station ID, collection date, collection position (lat then long), and sample depth. Each column header indicates what information is in each column.

The following parameters (units in brackets) are reported in this dataset: Date of collection, Latitude of collection (Decimal Degrees N), Longitude of collection (Decimal Degrees E), Sample Depth (m), ^{234}Th (dpm L^{-1}), ^{234}Th error (dpm L^{-1}), ^{238}U (dpm L^{-1}), ^{238}U error (dpm L^{-1}).

No data flags were used in the dataset.

This is data version 1 and was compiled on 10 May 2011, and is classified as FINAL.

DATA REMARKS:

None

---REFERENCES:

Buessler et al. 1992. Determination of thorium isotopes in seawater by non-destructive and radiochemical procedures. *Deep Sea Research I*. 39 (7/8): 1103-1114.

Buessler et al. 2001. An intercomparison of small- and large-volume techniques for thorium-234 in seawater. *Marine Chemistry*. 74:15-28.

Chen et al. 1986. ^{238}U , ^{234}U , and ^{232}Th in seawater. *Earth and Planetary Science Letters*. 80: 241-251.

