

Technical Report #08012 from
Bigelow Laboratory for Ocean Sciences

NUTRIENT, CHLOROPHYLL, and HYDROGRAPHIC PROFILES
at the NORTH POLE ENVIRONMENTAL OBSERVATORY,
April 18-28, 2005

Principal
Investigators:

Dr. J.P. Christensen
Bigelow Laboratory
180 McKown Pt.
W. Boothbay Hbr ME 04575
207-633-9600
jchristensen @ bigelow.org



Dr. W.S. Pegau
Dept. Oceanic Atmos. Sci.
104 Ocean Admin. Bldg.
Oregon State Univ.
Corvallis OR 97331

Grant Title: Collaborative Research: Annual Cycles of Nitrate & Phytoplankton Stocks
using optics at the North Pole Environmental Observatory
(OPP-0352641 to Christensen, OPP0352656 to Pegau)

The goal of this project was to begin an evaluation of nutrients and phytoplankton standing stocks in the central Arctic Ocean based on sampling at the North Pole Environmental Observatory. The first sampling occurred in April 2005 where eight vertical profiles of hydrographic properties (salinity, temperature, sigma-t), in-situ fluorescence, and nutrients in the region of the NPEO were determined (Figure 1). Station locations, times, and sampling details are tabulated in file, NPB05VIT.CSV, and column names and their descriptions in Appendix 1 of this report.

Vertical profiles of hydrographic, fluorescence, and nutrients over the upper 150 m were obtained using a Seabird SBE-19 CTD (serial number 1923347-2995) combined with a Wetstar model WS3S miniature flow-through fluorometer (serial number 646P), which had an excitation wavelength of 460 nm and an emission wavelength of 695 nm. The continuous CTD profiles were processed using standard SeaSoft software with the results being binned into 0.5 m depth intervals. The fluorometer was calibrated in the laboratory with aqueous solutions of resorufin (Christensen, 2006a). The fluorescence of resorufin was compared with the fluorescence of purified chlorophyll-a (from the algae *Anacystis nidulans*) dissolved in buffered acetone (10 ml of 1% (weight per volume) magnesium carbonate per L of 90% acetone). These chlorophyll solutions showed a fluorescence of 40.0 times that of resorufin for the same molar concentrations. The output of the fluorometer was first converted to fluorescence of resorufin ($\mu\text{mol/L}$) and then to that of chlorophyll-a (expressed in microgram-chl/L) using this factor and the molecular weight of chlorophyll-a (893.5 g/mole). The continuous vertical profiles for all stations are tabulated in the file, NPB05CTD.CSV, and column names and their descriptions in Appendix 2 of this report.

Figure 1. Locations sampled during NPEO 2005 and showing previous years' locations.



Nutrients and $^{18}\text{O} / ^{16}\text{O}$ ratios of the seawater were determined on samples collected with Niskin water bottles at 8-9 depths over the profile distance. These samples were frozen in tightly sealing bottles for return to Maine for measurement. Nutrients were measured by autoanalyzer. Nitrate and nitrite concentrations were measured by the methods of Armstrong et al. (1967) and Pavlou (1972). Ammonium was measured using the method of Koroleff (1970) and Slawyk and Maclsaac (1972). Phosphate was measured by the method of Murphy and Riley (1962) and Pavlou (1972). Dissolved silicate was measured by the method of Armstrong et al. (1967). The oxygen isotope data was measured at the Stable Isotope Facility at the University of Maine at Orono. The discrete vertical profiles for all stations are tabulated in the file, NPB05BTL.CSV, and column names and their descriptions in Appendix 3 of this report.

REFERENCES

Armstrong, F.A.J., C.R. Stearns, J.D.H. Strickland. 1967. The measurement of upwelling and subsequent biological processes by means of the Technicon autoanalyzer and associated equipment. *Deep-Sea Research* 14: 381-389.

Christensen, J.P. 2006a. Chlorophyll Fluorometer calibration using resorufin as standard. Technical report # 06001 from Bigelow Laboratory for Ocean Sciences. Bigelow Laboratory, W. Boothbay Harbor, Maine, 04575, 6 pp.

Koroleff, F. 1970. Revised version of "Direct determination of ammonia in natural waters as indophenol blue, *Int. Con. Explor. Sea, C.M. 1969/C:9*", ICES Information on Techniques and Methods for Sea Water Analysis, Interlab. Rep. No. 3, 19-22.

Murphy, J., J.P. Riley. 1962. A modified single solution method for the determination of phosphate in natural waters. *Analytica Chimica Acta* 12: 162-176.

Slawyk, G., J.J. Maclsaac. 1972. Comparison of two automated ammonium methods in a region of coastal upwelling. *Deep-Sea Research* 19: 521-523.

Pavlou, S.P. 1972. Phytoplankton Growth Dynamics. Technical Series 1, Chemostat Methodology and Chemical Analyses. Special Report No. 52, Dept. Oceanography, University of Washington, Seattle, WA 98195, 130 pp.

Appendix 1. Column listings in the data file NPB05VIT.CSV. The file is an ASC-II comma-separated-value file. Header-3 are headers found in row 8. Header-4 are headers found in row 9. Header-5 are headers found in row 10. Columns with alphanumeric data have blank values of "--", while columns of numerical data have blank values of -9, -99, or -999. All Borneo CTD samplings occurred through ice-holes within 5 m of each other and this site was about 100 m from the airstrip.

Column #	Header-3	Header-4	Header-5	Description
1	CRUISE#	NP-	#	Cruise number: year in units position - cruise identifier in tenths place
2	STATION	--	#	Sequential biological station number
3	NAME	SITE	NAME	Site name or location
4	TASK	--	NOTE	Site task
5				Section Separator
6	YEAR	GMT	YYYY	Sampling year at the station beginning
7	MONTH	GMT	MM	Sampling month at the station beginning
8	DAY	GMT	DD	Sampling day at the station beginning
9	HR	GMT	HH	Sampling hour at the station beginning
10	MIN	GMT	MM	Sampling minute at the station beginning
11	SEC	GMT	SS	Sampling second at the station beginning
12	JDAY	GMT	DECDAY	Julian day and time in decimal days at the station beginning
13	LAT	NORTH	DECDEG	North Latitude in decimal degrees at the station beginning
14	LONG	EAST	DECDEG	East Longitude in decimal degrees at the station beginning
15				Section Separator
16	DAY	GMT	DD	Sampling day at the station end
17	HR	GMT	HH	Sampling hour at the station end
18	MIN	GMT	MM	Sampling minute at the station end
19	SEC	GMT	SS	Sampling second at the station end
20	JDAY	GMT	DECDAY	Julian day and time in decimal days at the station end
21	LAT	NORTH	DECDEG	North latitude in decimal degrees at the station end
22	LONG	EAST	DECDEG	East longitude in decimal degrees at the station end
23				Section Separator
24	DISTANCE	FROM CTD1	KM	Distance from the CTD location of CTD-1 (beginning locations)
25	DRIFTRATE	LAST CTD	CM/S	Rate of drift between the current CTD cast and the previous CTD cast using beginning positions
26	DRIFTRATE	DURING CAST	CM/S	Rate of drift between the beginning and end of the current CTD cast
27	ICE	THICKNESS	CM	Thickness of ice-pack at the station
28	DEPTH	CTDMAX	M	Maximum depth of the CTD cast
29	NOTES	--	NOTE	Notes about the station

Appendix 2. Column listings in the data file NPB05CTD.CSV. The file is an ASC-II comma-separated-value file. Columns with alphanumeric data have blank values of "--", while columns of numerical data have blank values of -9, -99, or -999. Header-3 are headers found in row 8. Header-4 are headers found in row 9. Header-5 are headers found in row 10.

Column#	Header-3	Header-4	Header-5	Description
1	CRUISE#	--	#	Cruise number: year in units position - cruise identification in tenths place
2	CTD#	--	#	Sequential CTD number
3	DEPTH-BIN	BIN-CENTER	M	Center of the depth bin in meters
4	DEPTH-AV	--	M	Average depth of binned data in meters
5	PRES	--	DBAR	Average pressure of the binned data in decibars
6	TEMP	--	DEGC	Average temperature of the binned data in degrees C
7	SAL	--	PSU	Average salinity of the binned data in psu
8	SIGMAT	--	MG/L	Average density (sigma-t) of the binned data in mg/liter
9	FLUOR	[CHLA]EX	UG-CHL/L	Average fluorometer-assessed chlorophyll concentration in micrograms of extracted chlorophyll-a per liter

Appendix 3. Column listings in the data file NPB05BTL.CSV. The file is an ASC-II comma-separated-value file. Columns with alphanumeric data have blank values of "--", while columns of numerical data have blank values of -9, -99, or -999. Header-3 are headers found in row 8. Header-4 are headers found in row 9. Header-5 are headers found in row 10.

Column #	Header-3	Header-4	Header-5	DESCRIPTION
1	CRUISE#	NP-	#	Cruise number: year in units position - cruise identifier in tenths place
2	STATION	--	#	Sequential biological station #
3	NAME	SITE	NAME	Site name or location
4	DEPTH-BIN	BIN-CENTER	M	Selected bottle depth in meters
5	DEPTH-BTL	--	M	CTD-measured bottle depths in meters
6	PRES	--	DBAR	CTD-measured pressures in decibars
7	TEMP	--	DEGC	CTD-measured temperatures in degrees C
8	SAL	--	PSU	CTD-measured salinities in psu
9	SIGMAT	--	MG/L	Density (sigma-t) values in mg/liter
10	FLUOR	[CHLA]EX	UG-CHL/L	In-situ fluorometer measured chlorophyll concentrations in micrograms of extracted chlorophyll-a per liter
11	NO3	--	UMOL/L	Nitrate concentrations in micromoles/L
12	NO2	--	UMOL/L	Nitrite concentrations in micromoles/L
13	NH4	--	UMOL/L	Ammonium concentrations in micromoles/L
14	TIN	--	UMOL/L	Total inorganic nitrogen concentrations (sum of nitrate, nitrite, ammonium) in micromoles/L
15	PO4	--	UMOL/L	inorganic phosphate concentrations in micromoles/L
16	SI	--	UMOL/L	dissolved inorganic silicate concentrations in micromoles/L
17	O18/O16	--	PER MIL	seawater ¹⁸ O isotope ratios in per mil relative to SMOW

Report Date: 12 April 2008
 Report End: file NPB05DOC.XXX