

## Title:

BE CO<sub>2</sub> Flux Summer 2009

## Authors:

Walter C. Oechel and Cove S. Sturtevant

Global Change Research Group

Department of Biology, PS 241E

5500 Campanile Dr.

San Diego, CA 92182

Phone: 1-619-594-6613

Fax: 1-619-594-7831

PI Email: [oechel@sunstroke.sdsu.edu](mailto:oechel@sunstroke.sdsu.edu)

Data Questions Email: [sturteva@sciences.sdsu.edu](mailto:sturteva@sciences.sdsu.edu)

WWW address: <http://gcrp.sdsu.edu>

## Funding Source:

This research was conducted as part of the Biocomplexity Program of the National Science Foundation, grant number OPP-0421588.

## Data Set Overview:

This data set contains half-hourly vertical fluxes of CO<sub>2</sub> and related environmental measurements taken at the Biocomplexity Experimental Site in northern Alaska during the summer of 2009. Sampling was conducted as part of a large-scale water manipulation experiment to test the effect of altered tundra moisture on ecosystem carbon exchange.

The variables measured were: CO<sub>2</sub> flux H<sub>2</sub>O flux, latent and sensible heat fluxes, turbulence characteristics, radiation, meteorological variables, ground heat flux (5 replicates), and soil temperature (1 cm depth). Sample date and time are also given.

## Citing These Data:

Oechel, W.C., C.S. Sturtevant. 2011. *Carbon dioxide fluxes at the Biocomplexity Experiment Site, Barrow, AK, 2009*. Boulder, CO: National Center for Atmospheric Research, ARCSS Data Archive.

## Overview Table:

Category	Description
<a href="#">Data format</a>	Comma-delimited ASCII text
<a href="#">Spatial coverage and resolution</a>	Biocomplexity Experiment Site near Barrow, Alaska at 71.284° N, 156.598° W. Three meteorological towers spaced approximately 300 m apart in a North-South direction.
<a href="#">Temporal coverage and resolution</a>	2009-06-01 to 2009-09-01; measurements recorded in half-hourly intervals
<a href="#">File size</a>	3 files. BECO2FluxSummer2009_North.csv = 721 KB BECO2FluxSummer2009_Central.csv = 609 KB

	BECO2FluxSummer2009_South.csv = 721 KB
<a href="#">Parameter(s)</a>	Fc, H, FH2O,LE, TAU, Ta, Rh, VPD, Rn, Rgs (BEC only), PAR, WS, WD, UST, ZL, PRESS, PREC (BEC only),FG1, FG2, FG3, FG4, FG5, TS1

## Table of Contents

- [1. Contacts and Acknowledgments](#)
- [2. Detailed Data Description](#)
- [3. Data Access and Tools](#)
- [4. Data Acquisition and Processing](#)
- [5. References](#)
- [6. Document Information](#)

## 1. Contacts and Acknowledgments:

### Investigator Name and Title:

Walter C. Oechel  
Global Change Research Group  
Department of Biology  
San Diego State University  
5500 Campanile Ave.  
San Diego, CA, USA 92182

### Technical Contact:

Cove S. Sturtevant  
Global Change Research Group  
Department of Biology  
San Diego State University  
5500 Campanile Ave.  
San Diego, CA, USA 92182

## 2. Detailed Data Description:

### Format:

The data are available as comma-delimited ASCII text files. Separate files contain data from each section (North, Central, South), and are labeled accordingly. In each file, a 5-row header indicates:

Site name: <BEN = North (flooded), BEC = Central (drained), BES = South (intermediate)>

Contact email

Date of file creation

Column variable names

Column variable units

### File Size:

The file size for each file is as follows:

BECO2FluxSummer2009\_North.csv = 721 KB

BECO2FluxSummer2009\_Central.csv = 609 KB

BECO2FluxSummer2009\_South.csv = 721 KB

### **Spatial Coverage:**

Data were collected at the Biocomplexity Experimental Site in northern Alaska, located at 71.284° N, 156.598° W.

The Biocomplexity Experiment site is situated in the Barrow Environmental Observatory and located approximately 6 km east of Barrow village and 7 km south of the Arctic Ocean. The tundra in this region is considered wet-sedge meadow and is underlain by continuous permafrost with an average active layer of approximately 37 cm. Soils for this site are Gelisols of the suborder Histel. Mean annual temperature is -12°C, while mean summer temperature (June-August) is 3.3°C. Annual precipitation averages 106 mm.

The Biocomplexity Site is an approximately 1.5 km x 0.5 km vegetated thaw lake basin which is estimated to have drained 50-300 years ago. This basin has been divided into three sections by dikes. In 2009, water was pumped into or out of each section such that the North section represented wetter than ambient conditions, the Central section drier than ambient conditions, and the South section intermediate moisture conditions. The sections are about 300 m apart in the north-south direction. This was the second successful year of manipulation.

### **Spatial Resolution:**

All reported measurements were collected at the eddy-covariance tower located in the east of each section. Data from each section are contained in separate files.

### **Temporal Coverage:**

Data were collected from June 1, 2009 to September 1, 2009 (Julian days 152 - 244).

### **Temporal Resolution:**

All measurements are reported in half-hour intervals, with the indicated time of measurement as the beginning of the half-hour.

### **Parameters or Variables:**

#### **Parameter Description:**

Year: Year of measurement.

DOY: Day of year of measurement.

HRMIN: Hour and minute of measurement given as a 2 to 4 value integer, with the hour indicated in the first two values and the minute in the second two values. Ex. 12:30am is represented as 30, 8:30am is represented as 830, and 6:00pm is represented as 1800.

DTIME: Decimal day of year.

Fc: CO<sub>2</sub> flux in  $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ .

H: Sensible heat flux in  $\text{W m}^{-2}$

FH<sub>2</sub>O: Water vapor flux in  $\text{mmol H}_2\text{O m}^{-2} \text{ s}^{-1}$

LE: Latent heat flux in  $\text{W m}^{-2} \text{ s}^{-1}$

TAU: Momentum flux in  $\text{kg m}^{-1} \text{ s}^{-2}$

Ta: Air temperature at 1.6 m height in °C

Rh: Relative humidity at 1.6 m height in %

VPD: Vapor pressure deficit in kPa

Rn: Net radiation in  $\text{W m}^{-2}$

Rgs (BEC only): Downwelling global shortwave radiation (310 – 2800 nm)

PAR: Downwelling photosynthetically active radiation (400 – 700 nm)

WS: Wind speed in  $\text{m s}^{-1}$

WD: Wind direction in degrees clockwise from North

UST: Friction velocity in  $\text{m s}^{-1}$

ZL: Stability parameter (measurement height divided by the Obukhov Length) in  $\text{m m}^{-1}$

PRESS: Atmospheric pressure in kPa

PREC (BEC only): Precipitation summed for period of record in mm

FG1: Soil heat flux at 2 cm below surface in  $\text{W m}^{-2}$

FG2: Soil heat flux at 2 cm below surface in  $\text{W m}^{-2}$

FG3: Soil heat flux at 2 cm below surface in  $\text{W m}^{-2}$

FG4: Soil heat flux at 2 cm below surface in  $\text{W m}^{-2}$

FG5: Soil heat flux at 2 cm below surface in  $\text{W m}^{-2}$

TS1: Soil temperature at 1 cm depth in degrees Celsius

Missing/bad records are indicated with a -9999 value. Positive flux values (Fc, FH<sub>2</sub>O, H, LE, TAU) indicate upward fluxes. Positive radiation values (Rn, Rgs, PAR, FG) indicate downward fluxes.

### Sample Data Record:

YEA R	DO Y	HRMIN	DTIME	Fc umol CO2 m-2 s-1	H W m-2	FH2O mmol H2O m-2 s-1	LE W m-2	TAU kg m-1 s-2	Ta deg C	Rh %	VP D kP a
2009	152	0	152.01	0.1066	-15.742	-0.16467	-7.4314	0.024181	-2.6154	91.6	42
2009	152	30	152.03	0.15618	-17.622	-0.19859	-8.9662	0.029578	-3.1084	91.7	4
2009	152	100	152.05	0.13727	-17.702	-0.18134	-8.1879	0.026159	-3.1588	91.7	4
2009	152	130	152.07	0.055067	-11.139	-0.11977	-5.4115	0.013309	-3.8546	91.2	41
2009	152	200	152.09	-9999	-9999	-9999	-9999	0.0053569	-4.3188	91	4
2009	152	230	152.11	-9999	-9999	-9999	-9999	0.011322	-4.4448	92.2	34
2009	152	300	152.14	-9999	-9999	-9999	-9999	0.010999	-4.2681	91.4	38

## 3. Data Access and Tools:

### Data Access:

Data are available for ordering from [NCAR](#).

## 4. Data Acquisition and Processing:

### Sensor or Instrument Description:

Fluxes were measured at a height of 1.4 m above the terrain using traditional eddy covariance instrumentation. Flux data were taken at 10 Hz and recorded with a datalogger. Three-dimensional wind speed and virtual sonic temperature were collected with a Gill Wind Master Pro sonic anemometer (Gill Instruments Ltd., Hampshire, England). Molar densities of CO<sub>2</sub> and H<sub>2</sub>O were collected with an open path LI-7500 infrared gas analyzer (LI-COR Biosciences, Lincoln, NE, USA).

The LI-7500 gas analyzers were calibrated via a 2-point linear equation every 4 weeks. Ultra high purity nitrogen was used as the zero for CO<sub>2</sub> and H<sub>2</sub>O. High precision gas with certification accuracy < 1% was used as a span value for CO<sub>2</sub> (750 ppm in a balance of nitrogen). The span value for H<sub>2</sub>O was generated with a LI-610 dew point generator (LI-COR Biosciences, Lincoln, NE, USA) using a dew point 5 °C below ambient air temperature.

Environmental measurements were recorded in the immediate vicinity of each tower and within the manipulation footprint. Air temperature and relative humidity were measured at 1.6 m height with a HMP45C probe with radiation shield (Vaisala, Helsinki, Finland). Atmospheric pressure was measured with the LI-7500. Temperatures at 1 cm soil depth were measured with type T thermocouples. A 1.5 m tripod housed radiation instruments measuring: downwelling global shortwave radiation (BEC only; 310 – 2800 nm) with CMP3 pyranometers (Kipp and Zonen, Delft, The Netherlands), downwelling photosynthetically-active radiation (PAR, 400-700 nm) with LI-190 quantum sensors (LI-COR Biosciences, Lincoln, NE, USA), and net radiation (0.25 – 60  $\mu\text{m}$ ) with a Q7 net radiometer (REBS, Bellvue, Washington, USA). Ground heat flux at 2 cm depth was recorded with five HFT3 ground heat flux plates (REBS, Bellvue, Washington, USA). Finally, a TR-525M tipping rain gauge bucket measured liquid precipitation (Texas Electronics, Dallas, Texas, USA) at the Central section. These environmental variables were measured at 1 Hz and averaged into half-hour blocks.

### **Processing:**

Half-hourly flux calculations of carbon dioxide, water vapour, energy, and momentum were made using the eddy covariance method (Baldocchi et al. 1988) and coded in MATLAB v. 7.2 (Mathworks, Natick, Massachusetts, USA). Prior to covariance computations for each half hour, the sonic anemometer coordinate frame was double-rotated to align with the mean streamline and signals from separate sensors were time-aligned by maximizing the cross-correlation between vertical wind speed and scalar concentration. Appropriate corrections were applied for the simultaneous vertical transfer of heat and water vapour (Webb et al. 1980) as well as for high frequency spectral loss due to sensor separation and path length averaging (Moore 1986). No gap-filling was performed.

Quality control was applied pre- and post- flux computation. Raw data were de-spiked ( $> 6$  standard deviations from the running mean) and removed of periods clearly demonstrating error due to heavy mist, rain, or snow. Computed fluxes were filtered according to stationarity and integral turbulence characteristics tests following Foken et al. (2004). Fluxes calculated under low turbulence conditions (friction velocity  $< 0.1$  m/s) or when winds flowed from outside the manipulation footprint were excluded.

## **5. References and Related Publications**

### **References:**

- Baldocchi DD, Hicks BB, Meyers TP. 1988. MEASURING BIOSPHERE-ATMOSPHERE EXCHANGES OF BIOLOGICALLY RELATED GASES WITH MICROMETEOROLOGICAL METHODS. *Ecology* 69(5):1331-1340.
- Foken T, Gockede M, Mauder M *et al*: Post-field data quality control. In: *Handbook of micrometeorology: a guide for surface flux measurement and analysis*. Edited by Lee X, Massman WJ, Law BE. Dordrecht: Kluwer Academic Publishers; 2004.
- Moore CJ. 1986. FREQUENCY-RESPONSE CORRECTIONS FOR EDDY-CORRELATION SYSTEMS. *Boundary-Layer Meteorology* 37(1-2):17-35.
- Webb EK, Pearman GI, Leuning R. 1980. CORRECTION OF FLUX MEASUREMENTS FOR DENSITY EFFECTS DUE TO HEAT AND WATER-VAPOR TRANSFER. *Quarterly Journal of the Royal Meteorological Society* 106(447):85-100.

**Related Publications:**

Sturtevant, CS and W Oechel. Submitted. Soil moisture control over autumn season methane flux, Arctic Coastal Plain of Alaska.

Zona D, Oechel WC, Kochendorfer J, Paw U KT, Salyuk AN, Olivas PC, Oberbauer SF, and Lipson DA. 2009. Methne fluxes during the initiation of a large-scale water table manipulation experiment in the Alaskan Arctic tundra. *Global Biogeochemical Cycles* 23. GB2013. Doi: 10.1029/2009GB003487.

**6. Document Information:****Document Creation Date:**

2011-06-23