

TITLE

CAMP_Tibet_BJ-Tower_20021001_20030331.stm

CONTACT

Hirohiko Ishikawa
Disaster Prevention Research Institute, Kyoto University
okasho, Uji, Kyoto Pref.,611-0011 Japan
Phone: +81-774-38-4159
Fax : +81-774-38-4158
Email: ishikawa@storm.dpri.kyoto-u.ac.jp

Ken'ich UENO
University of Shiga Prefecture
Hassaka 2500 Hikone, Shiga 522-8533, Japan
Phone: +81-749-28-8312
Fax : +81-749-28-8477
Email: kueno@ses.usp.ac.jp

Yaoming MA
Institute for Tibetan Plateau Research
P.O. Box 2871, Beijing 100085, China.
Phone: +86-10-6284-9294
Fax : +86-10-6284-9886
Email: ymma@itp.cas.ac.cn

Kenji Tanaka
Department of Civil and Environmental Engineering, Kumamoto University
Kurokami 2-39-1, Kumamoto, Kumamoto Pref., 860-8555, Japan
Phone: +81-96-342-3601
Fax : +81-96-342-3601
Email: ktanaka@gpo.kumamoto-u.ac.jp

DATE OF THIS DOCUMENT

02 Sep. 2004 (Updated 31 Aug. 2006)

1. 0 DATASET OVERVIEW

1.1 Introduction

To clarify the energy and water cycle in the Tibetan Plateau, it is important to understand the characteristics of the basic meteorological elements and surface fluxes.

The purpose of Tibet AWS (Automatic Weather Station) observation is to improve the quantitative understanding of land-atmosphere interactions over the Tibetan Plateau and develop the land surface process models by monitoring these meteorological values.

1.2 Time period covered by the data

Start: 1 October 2002, 00:00

End: 31 March 2003, 23:00

1.3 Temporal characteristics of the data

All parameters are recorded every hour.

1.4 Physical location of the measurement

Latitude : 31.36866 N

Longitude : 91.89871 E

Elevation : 4509.2 m a.s.l.

Landscape : Bare land (with the thin weed-like plant)

Canopy height : Less than 5cm.

Soil Characteristics: Sand

1.5 Data source

1.6 Website address references

<http://monsoon.t.u-tokyo.ac.jp/camp/tibets/>

2.0 INSTRUMENTATION DESCRIPTION

2.1 Platform

The BJ site is located about 20 km southwest from the city of Naqu in the eastern Tibetan Plateau. This AWS was constructed in summer 2000. The system is originally same as that of D105, MS3478 (N-PAM) and ANNI. The direct incoming solar radiation and the scattered solar radiation measurement were added in June 2002. BJ site is the most enhanced observation site in the Tibetan Plateau: not only hydro-meteorological observation by AWS and SMTMS system, but atmospheric profile measurement by wind profiler, radiosonde, etc.

2.2 Description of the instrumentation

Parameter	Model	Manufacturer
Soil Temperature	Pt100	VAISALA

2.3 Instrumentation specification

Soil Temp_0cm : Soil Temperature at the 0cm depth (deg.C)

Soil Temp_4cm : Soil Temperature at the 4cm depth (deg.C)

Soil Temp_10cm : Soil Temperature at the 10cm depth (deg.C)

Soil Temp_20cm : Soil Temperature at the 20m depth (deg.C)

Soil Temp_40cm : Soil Temperature at the 40m depth (deg.C)

3.0 DATA COLLECTION AND PROCESSING

3.1 Description of data collection

Original data are sampled at every 5 seconds (0.2Hz) and 10-minute average is computed and stored in a data logger (Campbell CR-10X).

Data are downloaded from the Tower twice every year, in spring and summer. Then, data are sent to Japan, where they are processed.

3.2 Description of derived parameters and processing techniques used

Soil temperature is averaged over the previous hour.

Soil Moisture instantaneous values of each 1 hour.

There are two Soil Temperature sensors at the 0 cm depth. This time we apply the average of these two data to get the representative value at 0 cm.

4.0 QUALITY CONTROL PROCEDURES

For all parameters, the data has been visually checked, looking for extremely and unusual low/high values and/or periods with constant values thorough the CAMP Quality Control Web Interface.

The quality control flags follow the CEOP data flag definition document.

5.0 GAP FILLING PROCEDURES

No gap filling procedure was applied.

6.0 DATA REMARKS

6.1 PI's assessment of the data

6.1.1 Instruments problems

None

6.1.2 Quality issues

6.2 Missing data periods

None

7.0 REFERENCE REQUIREMENTS

Original data was collected and is provided within the framework of GAME/CAMP Tibet Scientific and Technological Research Project, funded by the Ministry of Education, Culture, Sports, Science and Technology; the Japan Science and Technology Agency; the Frontier Research System for Global Change; the Japan Aerospace Exploration Agency; the Chinese Academy of Sciences; and the Chinese Academy of Meteorological Sciences.

8.0 REFERENCES

H. Ishikawa and GAME-Tibet Boundary Layer Group, 2001: What has been known and what has not in GAME/Tibet BL observation, Proceedings of the Fifth International Study Conference on GEWEX in Asia and GAME, 691.

Ma, Yaoming, O. Tsukamoto, H. Ishikawa, Z. Su, M. Menenti, J. Wang and J. Wen, 2002: Determination of regional land surface heat flux densities over heterogeneous landscape of HEIFE integrating satellite remote sensing with field observations, Jour. Meteorol. Soc. Japan, 80(3), 485-501.

K. Tanaka, I. Tamagawa, H. Ishikawa, Y. Ma and Z. Hu, 2003: Surface energy and closure of the eastern Tibetan Plateau during the GAME-Tibet IOP 1998, J. Hydrology, vol. 283, pp. 169-183

K. Tanaka and H. Ishikawa, 2001: Long term monitoring of surface energy fluxes of the Amdo PBL site in the eastern Tibetan Plateau, Proceedings of the Fifth International Study Conference on GEWEX in Asia and GAME, 384-388.

Ueno, K., H. Fujii, H. Yamada and L. Liu, (2001) Weak and Frequent Monsoon Precipitation over the Tibetan Plateau. J. Meteor. Soc. Japan, 79, 1B, 419-434.

TITLE

CAMP_Tibet_BJ-Tower_20030401_20030930.stm

CONTACT

Hirohiko Ishikawa
Disaster Prevention Research Institute, Kyoto University
Gokasho, Uji, Kyoto Pref.,611-0011 Japan
Phone: +81-774-38-4159
Fax : +81-774-38-4158
Email: ishikawa@storm.dpri.kyoto-u.ac.jp

Ken'ich UENO
University of Tsukuba
Tennoudai 1-1-1 Tsukuba, Ibaraki 305-8572, Japan
Phone/Fax: +8129-853-4399
Email: kenueno@sakura.cc.tsukuba.ac.jp

Yaoming MA
Institute for Tibetan Plateau Research
P.O. Box 2871, Beijing 100085, China.
Phone: +86-10-6284-9294
Fax : +86-10-6284-9886
Email: ymma@itp.cas.ac.cn

Kenji Tanaka
Department of Civil and Environmental Engineering, Kumamoto University
Kurokami 2-39-1, Kumamoto, Kumamoto Pref., 860-8555, Japan
Phone/Fax: +81-96-342-3601
Email: ktanaka@gpo.kumamoto-u.ac.jp

DATE OF THIS DOCUMENT

19 Apr. 2006 (**Updated 31 Aug. 2006**)

1. 0 DATASET OVERVIEW

1.7 Introduction

To clarify the energy and water cycle in the Tibetan Plateau, it is important to understand the characteristics of the basic meteorological elements and surface fluxes.

The purpose of Tibet AWS (Automatic Weather Station) observation is to improve the quantitative understanding of land-atmosphere interactions over the Tibetan Plateau and develop the land surface process models by monitoring these meteorological values.

1.8 Time period covered by the data

Start: 1 April 2003, 00:00
End: 30 September 2003, 23:00

1.9 Temporal characteristics of the data

All parameters are recoded every hour.

1.10 Physical location of the measurement

Latitude : 31.36866 N
Longitude : 91.89871 E
Elevation : 4509.2 m a.s.l.
Landscape : Bare land (with the thin weed-like plant)
Canopy height : Less than 5cm.
Soil Characteristics: Sand

1.11 Data source

1.12 Website address references

<http://monsoon.t.u-tokyo.ac.jp/camp/tibets/>

2.0 INSTRUMENTATION DESCRIPTION

2.1 Platform

The BJ site is located about 20 km southwest from the city of Naqu in the eastern Tibetan Plateau. This AWS was constructed in summer 2000. The system is originally same as that of D105, MS3478 (N-PAM) and ANNI. The direct incoming solar radiation and the scattered solar radiation measurement were added in June 2002. BJ site is the most enhanced observation site in the Tibetan Plateau: not only hydro-meteorological observation by AWS and SMTMS system, but atmospheric profile measurement by wind profiler, radiosonde, etc.

2.2 Description of the instrumentation

Parameter	Model	Manufacturer
Soil Temperature	Pt100	VAISALA

2.4 Instrumentation specification

Soil Temp_0cm : Soil Temperature at the 0cm depth (deg.C)
Soil Temp_4cm : Soil Temperature at the 4cm depth (deg.C)
Soil Temp_10cm : Soil Temperature at the 10cm depth (deg.C)
Soil Temp_20cm : Soil Temperature at the 20m depth (deg.C)
Soil Temp_40cm : Soil Temperature at the 40m depth (deg.C)

3.0 DATA COLLECTION AND PROCESSING

3.1 Description of data collection

Original data are sampled at every 5 seconds (0.2Hz) and 10-minute average is computed and stored in a data logger (Campbell CR-10X).

Data are downloaded from the Tower twice every year, in spring and summer. Then, data are sent to Japan, where they are processed.

3.2 Description of derived parameters and processing techniques used

Soil temperature is averaged over the previous hour.

Soil Moisture instantaneous values of each 1 hour.

There are two Soil Temperature sensors at the 0 cm depth. This time we apply the average of these two data to get the representative value at 0 cm.

4.0 QUALITY CONTROL PROCEDURES

For all parameters, the data has been visually checked, looking for extremely and unusual low/high values and/or periods with constant values thorough the CAMP Quality Control Web Interface.

The quality control flags follow the CEOP data flag definition document.

5.0 GAP FILLING PROCEDURES

No gap filling procedure was applied.

6.0 DATA REMARKS

6.1 PI's assessment of the data

6.1.1 Instruments problems

None

6.1.2 Quality issues

6.2 Missing data periods

Please see the chapter 9.0.

7.0 REFERENCE REQUIREMENTS

Original data was collected and is provided within the framework of GAME/CAMP Tibet Scientific and Technological Research Project, funded by the Ministry of Education, Culture, Sports, Science and Technology; the Japan Science and Technology Agency; the Frontier Research System for Global Change; the Japan Aerospace Exploration Agency; the Chinese Academy of Sciences; and the Chinese Academy of Meteorological Sciences.

8.0 REFERENCES

H. Ishikawa and GAME-Tibet Boundary Layer Group, 2001: What has been known and what has not in GAME/Tibet BL observation, Proceedings of the Fifth International Study Conference on GEWEX in Asia and GAME, 691.

Ma, Yaoming, O. Tsukamoto, H. Ishikawa, Z. Su, M. Menenti, J. Wang and J. Wen, 2002: Determination of regional land surface heat flux densities over heterogeneous landscape of HEIFE integrating satellite remote sensing with field observations, Jour. Meteorol. Soc. Japan, 80(3), 485-501.

K. Tanaka, I. Tamagawa, H. Ishikawa, Y. Ma and Z. Hu, 2003: Surface energy and closure of the eastern Tibetan Plateau during the GAME-Tibet IOP 1998, J. Hydrology, vol. 283, pp. 169-183

K. Tanaka and H. Ishikawa, 2001: Long term monitoring of surface energy fluxes of the Amdo PBL site in the eastern Tibetan Plateau, Proceedings of the Fifth International Study Conference on GEWEX in Asia and GAME, 384-388.

Ueno, K., H. Fujii, H. Yamada and L. Liu, (2001) Weak and Frequent Monsoon Precipitation over the Tibetan Plateau. J. Meteor. Soc. Japan, 79, 1B, 419-434.

9.0 Missing data periods

None

TITLE

[CAMP_Tibet_BJ-Tower_20031001_20040830.stm](#)

CONTACT

Hirohiko Ishikawa
Disaster Prevention Research Institute, Kyoto University
Gokasho, Uji, Kyoto Pref.,611-0011 Japan
Phone: +81-774-38-4159
Fax : +81-774-38-4158
Email: ishikawa@storm.dpri.kyoto-u.ac.jp

Ken'ich UENO
University of Tsukuba
Tennoudai 1-1-1 Tsukuba, Ibaraki 305-8572, Japan
Phone/Fax: +8129-853-4399
Email: kenueno@sakura.cc.tsukuba.ac.jp

Yaoming MA
Institute for Tibetan Plateau Research
P.O. Box 2871, Beijing 100085, China.
Phone: +86-10-6284-9294
Fax : +86-10-6284-9886
Email: ymma@itp.cas.ac.cn

Kenji Tanaka
Department of Civil and Environmental Engineering, Kumamoto University
Kurokami 2-39-1, Kumamoto, Kumamoto Pref., 860-8555, Japan
Phone/Fax: +81-96-342-3601
Email: ktanaka@gpo.kumamoto-u.ac.jp

DATE OF THIS DOCUMENT

7 July, 2006

1. 0 DATASET OVERVIEW

1.13 Introduction

To clarify the energy and water cycle in the Tibetan Plateau, it is important to understand the characteristics of the basic meteorological elements and surface fluxes.

The purpose of Tibet AWS (Automatic Weather Station) observation is to improve the quantitative understanding of land-atmosphere interactions over the Tibetan Plateau and develop the land surface process models by monitoring these meteorological values.

1.14 Time period covered by the data

Start: 1 October 2003, 00:00
End: 30 August 2004, 23:00

1.15 Temporal characteristics of the data

All parameters are recoded every hour.

1.16 Physical location of the measurement

Latitude : 31.36866 N
Longitude : 91.89871 E
Elevation : 4509.2 m a.s.l.
Landscape : Bare land (with the thin weed-like plant)
Canopy height : Less than 5cm.
Soil Characteristics: Sand

1.17 Data source

1.18 Website address references

<http://monsoon.t.u-tokyo.ac.jp/camp/tibets/>

2.0 INSTRUMENTATION DESCRIPTION

2.1 Platform

The BJ site is located about 20 km southwest from the city of Naqu in the eastern Tibetan Plateau. This AWS was constructed in summer 2000. The system is originally same as that of D105, MS3478 (N-PAM) and ANNI. The direct incoming solar radiation and the scattered solar radiation measurement were added in June 2002. BJ site is the most enhanced observation site in the Tibetan Plateau: not only hydro-meteorological observation by AWS and SMTMS system, but atmospheric profile measurement by wind profiler, radiosonde, etc.

2.2 Description of the instrumentation

Parameter	Model	Manufacturer
Soil Temperature	Pt100	VAISALA
Soil Moisture	Trime EZ	IMKO

2.5 Instrumentation specification

Soil Temp_0cm : Soil Temperature at the 0cm depth (deg.C)
Soil Temp_4cm : Soil Temperature at the 4cm depth (deg.C)
Soil Temp_10cm : Soil Temperature at the 10cm depth (deg.C)
Soil Temp_20cm : Soil Temperature at the 20m depth (deg.C)
Soil Temp_40cm : Soil Temperature at the 40m depth (deg.C)
Soil Moist_4cm : Soil Moisture at the 4cm depth (%)
Soil Moist_20cm : Soil Moisture at the 20cm depth (%)

3.0 DATA COLLECTION AND PROCESSING

3.1 Description of data collection

Original data are sampled at every 5 seconds (0.2Hz) and 10-minute average is computed and stored in a data logger (Campbell CR-10X).

Data are downloaded from the Tower twice every year, in spring and summer. Then, data are sent to Japan, where they are processed.

3.2 Description of derived parameters and processing techniques used

Soil temperature is averaged over the previous hour.

Soil Moisture instantaneous values of each 1 hour.

There are two Soil Temperature sensors at the 0 cm depth. This time we apply the average of these two data to get the representative value at 0 cm.

4.0 QUALITY CONTROL PROCEDURES

For all parameters, the data has been visually checked, looking for extremely and unusual low/high values and/or periods with constant values through the CAMP Quality Control Web Interface.

The quality control flags follow the CEOP data flag definition document.

5.0 GAP FILLING PROCEDURES

No gap filling procedure was applied.

6.0 DATA REMARKS

6.1 PI's assessment of the data

6.1.1 Instruments problems

None

6.1.2 Quality issues

6.2 Missing data periods

Please see the chapter 9.0.

7.0 REFERENCE REQUIREMENTS

Original data was collected and is provided within the framework of GAME/CAMP Tibet Scientific and Technological Research Project, funded by the Ministry of Education, Culture, Sports, Science and Technology; the Japan Science and Technology Agency; the Frontier Research System for Global Change; the Japan Aerospace Exploration Agency; the Chinese Academy of Sciences; and the Chinese Academy of Meteorological Sciences.

8.0 REFERENCES

H. Ishikawa and GAME-Tibet Boundary Layer Group, 2001: What has been known and what has not in GAME/Tibet BL observation, Proceedings of the Fifth International Study Conference on GEWEX in Asia and GAME, 691.

Ma, Yaoming, O. Tsukamoto, H. Ishikawa, Z. Su, M. Menenti, J. Wang and J. Wen, 2002: Determination of regional land surface heat flux densities over heterogeneous landscape of HEIFE integrating satellite remote sensing with field observations, Jour. Meteorol. Soc. Japan, 80(3), 485-501.

K. Tanaka, I. Tamagawa, H. Ishikawa, Y. Ma and Z. Hu, 2003: Surface energy and closure of the eastern Tibetan Plateau during the GAME-Tibet IOP 1998, J. Hydrology, vol. 283, pp. 169-183

K. Tanaka and H. Ishikawa, 2001: Long term monitoring of surface energy fluxes of the Amdo PBL site in the eastern Tibetan Plateau, Proceedings of the Fifth International Study Conference on GEWEX in Asia and GAME, 384-388.

Ueno, K., H. Fujii, H. Yamada and L. Liu, (2001) Weak and Frequent Monsoon Precipitation over the Tibetan Plateau. J. Meteor. Soc. Japan, 79, 1B, 419-434.

9.0 Missing data periods

File Name : CAMP_Tibet_BJ-Tower_20031001_20040830.stm
Data Period : 2003/10/01 00:00 - 2004/08/30 23:00

Soil Temperature (-0.40m)

2004/04/25 04:00 - 2004/06/06 20:00 (1025)

2004/08/30 02:00 - 2004/08/30 23:00 (22)

Soil Temperature (-0.20m)

2004/04/25 04:00 - 2004/06/06 20:00 (1025)

2004/08/30 02:00 - 2004/08/30 23:00 (22)

Soil Temperature (-0.10m)

2004/04/25 04:00 - 2004/06/06 20:00 (1025)

2004/08/30 02:00 - 2004/08/30 23:00 (22)

Soil Temperature (-0.04m)

2004/04/25 04:00 - 2004/06/06 20:00 (1025)

2004/08/30 02:00 - 2004/08/30 23:00 (22)

Soil Temperature (0.00m)

2004/04/25 05:00 - 2004/06/06 21:00 (1025)

2004/08/30 02:00 - 2004/08/30 23:00 (22)

Soil Moisture (-0.40m)

2003/10/01 00:00 - 2004/08/30 23:00 (ALL)

Soil Moisture (-0.20m)

2003/10/01 00:00 - 2003/12/31 15:00 (2200)

2004/01/01 01:00 - 2004/02/08 08:00 (920)
2004/04/25 04:00 - 2004/06/06 20:00 (1025)
2004/08/30 02:00 - 2004/08/30 23:00 (22)

Soil Moisture (-0.10m)
2003/10/01 00:00 - 2004/08/30 23:00 (ALL)

Soil Moisture (-0.04m)
2003/10/01 00:00 - 2003/12/31 15:00 (2200)
2004/01/01 01:00 - 2004/02/08 08:00 (920)
2004/04/25 04:00 - 2004/06/06 20:00 (1025)
2004/08/30 02:00 - 2004/08/30 23:00 (22)

Soil Moisture (0.00m)
2003/10/01 00:00 - 2004/08/30 23:00 (ALL)