TITLE

CAMP_Tibet_ANNI-AWS_20030401_20030930.stm

CONTACT

Hirohiko Ishikawa

Disaster Prevention Research Institute, Kyoto University

Gokasho, Uji, Kyoto Pref.,611-0011 Japan

Phone: +81-774-38-4159 Fax: +81-774-38-4158

Email: ishikawa@storm.dpri.kyoto-u.ac.jp

Ken'ich UENO

University of Tsukuba

Tennoudai 1-1-1 Tsukuba, Ibaraki 305-8572, Japan

Phone/Fax: +8129-853-4399

Email: kenueno@sakura.cc.tsukuba.ac.jp

Yaoming MA

Institute for Tibetan Plateau Research P.O. Box 2871, Beijing 100085, China.

Phone: +86-10-6284-9294 Fax: +86-10-6284-9886 Email: ymma@itp.cas.ac.cn

Kenji Tanaka

Department of Civil and Environmental Engineering, Kumamoto University

Kurokami 2-39-1, Kumamoto, Kumamoto Pref., 860-8555, Japan

Phone/Fax: +81-96-342-3601

Email: ktanaka@gpo.kumamoto-u.ac.jp

DATE OF THIS DOCUMENT

19 Apr. 2006 (Updated 23 Jun. 06)

1. 0 DATASET OVERVIEW

1.1 Introduction

To clarify the energy and water cycle in the Tibetan Plateau, it is important to understand the characteristics of the basic meteorological elements and surface fluxes.

The purpose of Tibet AWS (Automatic Weather Station) observation is to improve the quantitative understanding of land-atmosphere interactions over the Tibetan Plateau and develop the land surface process models by monitoring these meteorological values.

1.2 Time period covered by the data

Start: 1 October 2002, 00:00 End: 30 September 2003, 23:00

1.3 Temporal characteristics of the data

All parameters are recoded every hour.

1.4 Physical location of the measurement

Latitude : 31.25442 N Longitude : 92.17241 E Elevation : 4480 m a.s.l.

Landscape : Bare land (with the thin weed-like plant): Plain land

Canopy height: Less than 5cm. Soil Characteristics: Silt loam

1.5 Data source

1.6 Website address references

http://monsoon.t.u-tokyo.ac.jp/camp/tibets/

2.0 INSTRUMENTATION DESCRIPTION

2.1 Platform

Three Small AWS systems are constructed in June 2002 to represent the local scale (100m ~ 1km) heterogeneity of the surface condition to measure surface and soil temperature, soil moisture and heat flux. The wind speed and direction are only measured as the meteorological element in this system. This S-AWS (No.1) is located about 340 m westward from main AWS of BJ site. The sensors are mounted on several heights.

2.2 Description of the instrumentation

Parameter	Model	Manufacturer
Soil Temperature	WST80-L5	AGEMATSU
Soil Moisture	Trime IT	IMKO

2.3 <u>Instrumentation specification</u>

Soil Temp_0cm : Soil Temperature at the 0cm depth (deg.C)
Soil Temp_4cm : Soil Temperature at the 4cm depth (deg.C)
Soil Temp_10cm : Soil Temperature at the 10cm depth (deg.C)
Soil Temp_20cm : Soil Temperature at the 20cm depth (deg.C)
Soil Temp_40cm : Soil Temperature at the 40cm depth (deg.C)
Soil Moist_4cm : Averaged Soil Moisture at 4cm depth (%)
Soil Moist_20cm : Averaged Soil Moisture at 20cm depth (%)

3.0 DATA COLLECTION AND PROCESSING

3.1 <u>Description of data collection</u>

Original data are sampled at every 5 seconds (0.2Hz) and 10-minute average (except soil moisture, sampled at every 10 minutes) is computed and stored in a data logger (Campbell CR-10X).

Data are downloaded from the Tower twice every year, in spring and summer. Then, data are sent to Japan, where they are processed.

3.2 <u>Description of derived parameters and processing techniques used</u>

Soil temperature is averaged over the previous hour. Soil Moisture instantaneous values of each 1 hour.

4.0 QUALITY CONTROL PROCEDURES

For all parameters, the data has been visually checked, looking for extremely and unusual low/high values and/or periods with constant values thorough the CAMP Quality Control Web Interface.

The quality control flags follow the CEOP data flag definition document.

5.0 GAP FILLING PROCEDURES

No gap filling procedure was applied.

6.0 DATA REMARKS

- 6.1 PI's assessment of the data
- 6.1.1 <u>Instruments problems</u>

None.

6.1.2 Quality issues

6.2 Missing data periods

Missing data periods were listed in the chapter 9.0.

7.0 REFERENCE REQUIREMENTS

Original data was collected and is provided within the framework of GAME/CAMP Tibet Scientific and Technological Research Project, funded by the Ministry of Education, Culture, Sports, Science and Technology; the Japan Science and Technology Agency; the Frontier Research System for Global Change; the Japan Aerospace Exploration Agency; the Chinese Academy of Sciences; and the Chinese Academy of Meteorological Sciences.

8.0 REFERENCES

H. Ishikawa and GAME-Tibet Boundary Layer Group, 2001: What has been known and what has not in GAME/Tibet BL observation, Proceedings of the Fifth International Study Conference on GEWEX in Asia and GAME, 691.

Ma, Yaoming, O. Tsukamoto, H. Ishikawa, Z. Su, M. Menenti, J. Wang and J. Wen, 2002: Determination of regional land surface heat flux densities over heterogeneous landscape of HEIFE integrating satellite remote sensing with field observations, Jour. Meteorol. Soc. Japan, 80(3), 485-501.

K. Tanaka, I. Tamagawa, H. Ishikawa, Y. Ma and Z. Hu, 2003: Surface energy and closure of the eastern Tibetan Plateau during the GAME-Tibet IOP 1998, J. Hydrology, vol. 283, pp. 169-183

K. Tanaka and H. Ishikawa, 2001: Long term monitoring of surface energy fluxes of the Amdo PBL site in the eastern Tibetan Plateau, Proceedings of the Fifth International Study Conference on GEWEX in Asia and GAME, 384-388.

Ueno, K., H. Fujii, H. Yamada and L. Liu, (2001) Weak and Frequent Monsoon Precipitation over the Tibetan Plateau. J. Meteor. Soc. Japan, 79, 1B, 419-434.

9.0 Missing data periods

Soil Temperature (-0.40m) 2003/04/01 00:00 - 2003/09/18 04:00 (4085)

Soil Temperature (-0.20m) 2003/04/01 00:00 - 2003/09/18 04:00 (4085)

Soil Temperature (-0.10m) 2003/04/01 00:00 - 2003/09/18 04:00 (4085)

Soil Temperature (-0.04m) 2003/04/01 00:00 - 2003/09/18 04:00 (4085)

Soil Temperature (0.00m) 2003/04/01 00:00 - 2003/09/18 04:00 (4085)

Soil Moisture (-0.20m) 2003/04/01 00:00 - 2003/09/18 04:00 (4085)

Soil Moisture (-0.04m) 2003/04/01 00:00 - 2003/09/18 04:00 (4085)

TITLE

CAMP Tibet ANNI-AWS 20031001 20040814.stm

CONTACT

Hirohiko Ishikawa

Disaster Prevention Research Institute, Kyoto University

Gokasho, Uji, Kyoto Pref.,611-0011 Japan

Phone: +81-774-38-4159 Fax: +81-774-38-4158

Email: ishikawa@storm.dpri.kyoto-u.ac.jp

Ken'ich UENO

University of Tsukuba

Tennoudai 1-1-1 Tsukuba, Ibaraki 305-8572, Japan

Phone/Fax: +8129-853-4399

Email: kenueno@sakura.cc.tsukuba.ac.jp

Yaoming MA

Institute for Tibetan Plateau Research P.O. Box 2871, Beijing 100085, China.

Phone: +86-10-6284-9294 Fax: +86-10-6284-9886 Email: ymma@itp.cas.ac.cn

Kenji Tanaka

Department of Civil and Environmental Engineering, Kumamoto University

Kurokami 2-39-1, Kumamoto, Kumamoto Pref., 860-8555, Japan

Phone/Fax: +81-96-342-3601

Email: ktanaka@gpo.kumamoto-u.ac.jp

DATE OF THIS DOCUMENT

7 July. 2006

1. 0 DATASET OVERVIEW

1.7 Introduction

To clarify the energy and water cycle in the Tibetan Plateau, it is important to understand the characteristics of the basic meteorological elements and surface fluxes.

The purpose of Tibet AWS (Automatic Weather Station) observation is to improve the quantitative understanding of land-atmosphere interactions over the Tibetan Plateau and develop the land surface process models by monitoring these meteorological values.

1.8 Time period covered by the data

Start: 1 October 2003, 00:00 End: 14 August 2004, 23:00

1.9 Temporal characteristics of the data

All parameters are recoded every hour.

1.10 Physical location of the measurement

Latitude : 31.25442 N Longitude : 92.17241 E Elevation : 4480 m a.s.l.

Landscape : Bare land (with the thin weed-like plant): Plain land

Canopy height : Less than 5cm. Soil Characteristics: Silt loam

1.11 Data source

1.12 Website address references

http://monsoon.t.u-tokyo.ac.jp/camp/tibets/

2.0 INSTRUMENTATION DESCRIPTION

2.1 Platform

Three Small AWS systems are constructed in June 2002 to represent the local scale (100m ~ 1km) heterogeneity of the surface condition to measure surface and soil temperature, soil moisture and heat flux. The wind speed and direction are only measured as the meteorological element in this system. This S-AWS (No.1) is located about 340 m westward from main AWS of BJ site. The sensors are mounted on several heights.

2.2 Description of the instrumentation

Parameter	Model	Manufacturer
Soil Temperature	WST80-L5	AGEMATSU
Soil Moisture	Trime IT	IMKO

2.4 <u>Instrumentation specification</u>

Soil Temp_0cm : Soil Temperature at the 0cm depth (deg.C)
Soil Temp_4cm : Soil Temperature at the 4cm depth (deg.C)
Soil Temp_10cm : Soil Temperature at the 10cm depth (deg.C)
Soil Temp_20cm : Soil Temperature at the 20cm depth (deg.C)
Soil Temp_40cm : Soil Temperature at the 40cm depth (deg.C)
Soil Moist_4cm : Averaged Soil Moisture at 4cm depth (%)
Soil Moist_20cm : Averaged Soil Moisture at 20cm depth (%)

3.0 DATA COLLECTION AND PROCESSING

3.1 <u>Description of data collection</u>

Original data are sampled at every 5 seconds (0.2Hz) and 10-minute average (except soil moisture, sampled at every 10 minutes) is computed and stored in a data logger (Campbell CR-10X).

Data are downloaded from the Tower twice every year, in spring and summer. Then, data are sent to Japan, where they are processed.

3.2 <u>Description of derived parameters and processing techniques used</u>

Soil temperature is averaged over the previous hour. Soil Moisture instantaneous values of each 1 hour.

4.0 QUALITY CONTROL PROCEDURES

For all parameters, the data has been visually checked, looking for extremely and unusual low/high values and/or periods with constant values thorough the CAMP Quality Control Web Interface.

The quality control flags follow the CEOP data flag definition document.

5.0 GAP FILLING PROCEDURES

No gap filling procedure was applied.

6.0 DATA REMARKS

- 6.1 PI's assessment of the data
- 6.1.1 <u>Instruments problems</u>

None.

6.1.2 Quality issues

6.2 Missing data periods

Missing data periods were listed in the chapter 9.0.

7.0 REFERENCE REQUIREMENTS

Original data was collected and is provided within the framework of GAME/CAMP Tibet Scientific and Technological Research Project, funded by the Ministry of Education, Culture, Sports, Science and Technology; the Japan Science and Technology Agency; the Frontier Research System for Global Change; the Japan Aerospace Exploration Agency; the Chinese Academy of Sciences; and the Chinese Academy of Meteorological Sciences.

8.0 REFERENCES

H. Ishikawa and GAME-Tibet Boundary Layer Group, 2001: What has been known and what has not in GAME/Tibet BL observation, Proceedings of the Fifth International Study Conference on GEWEX in Asia and GAME, 691.

Ma, Yaoming, O. Tsukamoto, H. Ishikawa, Z. Su, M. Menenti, J. Wang and J. Wen, 2002: Determination of regional land surface heat flux densities over heterogeneous landscape of HEIFE integrating satellite remote sensing with field observations, Jour. Meteorol. Soc. Japan, 80(3), 485-501.

K. Tanaka, I. Tamagawa, H. Ishikawa, Y. Ma and Z. Hu, 2003: Surface energy and closure of the eastern Tibetan Plateau during the GAME-Tibet IOP 1998, J. Hydrology, vol. 283, pp. 169-183

K. Tanaka and H. Ishikawa, 2001: Long term monitoring of surface energy fluxes of the Amdo PBL site in the eastern Tibetan Plateau, Proceedings of the Fifth International Study Conference on GEWEX in Asia and GAME, 384-388.

Ueno, K., H. Fujii, H. Yamada and L. Liu, (2001) Weak and Frequent Monsoon Precipitation over the Tibetan Plateau. J. Meteor. Soc. Japan, 79, 1B, 419-434.

9.0 Missing data periods

Soil Temperature (-0.40m) 2003/04/01 00:00 - 2003/09/18 04:00 (4085)

Soil Temperature (-0.20m) 2003/04/01 00:00 - 2003/09/18 04:00 (4085)

Soil Temperature (-0.10m) 2003/04/01 00:00 - 2003/09/18 04:00 (4085)

Soil Temperature (-0.04m) 2003/04/01 00:00 - 2003/09/18 04:00 (4085)

Soil Temperature (0.00m) 2003/04/01 00:00 - 2003/09/18 04:00 (4085)

Soil Moisture (-0.20m) 2003/04/01 00:00 - 2003/09/18 04:00 (4085)

Soil Moisture (-0.04m) 2003/04/01 00:00 - 2003/09/18 04:00 (4085)