#### TITLE:

CEOP\_Tsukuba\_NIAES-MASE\_20090101\_20090630.stm

# CONTACT(S):

1) Akira Miyata

National Institute for Agro-Environmental Sciences

Tsukuba 305-8604, Japan

E-mail: amiyat@niaes.affrc.go.jp

2) Masayoshi Mano

National Institute for Agro-Environmental Sciences

Tsukuba 305-8604, Japan

E-mail: mmano@niaes.affrc.go.jp

### **DATE OF THIS DOCUMENT:**

30 November 2009

#### 1. 0 DATASET OVERVIEW:

### 1.1 Introduction:

Mase paddy flux site was established in 1999 to monitor greenhouse gas exchange between paddy fields and the atmosphere, and since then, Mase site is operated as one of the key study sites of AsiaFlux (http://www.asiaflux.net/). Details of the study site and instrumentation are given in some references (Saito *et al.*, 2005; Miyata *et al.*, 2005; Han *et al.*, 2007; Saito *et al.*, 2007).

# 1.2 Time period covered by the data:

Start: 1 January 2009, 00:00 (UTC) End: 30 June 2009, 23:30 (UTC)

### 1.3 Physical location of the measurement:

Latitude: 36° 03′ 14.3″ N Longitude: 140° 01′ 36.9″ E

Elevation: 11 m a.s.l.

Landscape: Agricultural fields (paddy fields)

Soil characteristics: Soil type is Eutric Fluvisols. The site is flooded most of rice growing

season (from the beginning of May to mid-September).

#### 1.4 Data source:

Original data.

### 1.5 WWW address references:

http://www.asiaflux.net/network/007MSE 1.html

http://ecomdb.niaes.affrc.go.jp/e level page.php?select area=1045&select site=1121

#### 2.0 INSTRUMENTATION DESCRIPTION:

### 2.1 Platform:

Sensors are set around a 6-m tall mast on which meteorological sensors are mounted.

## 2.2 <u>Description of the instrumentation:</u>

Parameter	Model	Manufacturer	
Soil Temperature	-	Home-made	
Soil Temperature	-	Home-made	
Soil Temperature	-	Home-made	
Soil Temperature	-	Home-made	
Soil Temperature	-	Home-made	
Soil Temperature	-	Home-made	
Soil Moisture	TDR100	Campbell, Logan, UT, USA	
Soil Moisture	TDR100	Campbell, Logan, UT, USA	
Soil Moisture	TDR100	Campbell, Logan, UT, USA	
Soil Moisture	TDR100	Campbell, Logan, UT, USA	
Soil Moisture	TDR100	Campbell, Logan, UT, USA	

# 2.3 Instrumentation specification:

Parameter	Sensor Type	Depth of sensor (m)	Accuracy	Resolution
Soil Temperature	T-type thermocouple	0.01	-	-
Soil Temperature	T-type thermocouple	0.02 (0.025)		
Soil Temperature	T-type thermocouple	0.05	-	-
Soil Temperature	T-type thermocouple	0.10	-	-
Soil Temperature	T-type thermocouple	0.20	-	-
Soil Temperature	T-type thermocouple	0.40	-	-
Soil Moisture	Time domain	0.02 (0.025) 1	-	-
	reflectometry			
Soil Moisture	Time domain	0.03 (from surface to	-	-
	reflectometry	0.05) 2		
Soil Moisture	Time domain	0.05 (from surface to	-	-
	reflectometry	0.08) 2		
Soil Moisture	Time domain	0.10 (from surface to	-	-
	reflectometry	0.20) 2		
Soil Moisture	Time domain	0.15 (from surface to	-	-
	reflectometry	0.30) 2,3		

<sup>&</sup>lt;sup>1</sup> The sensor was set horizontally in the soil at 2.5 cm depth.

### 3.0 DATA COLLECTION AND PROCESSING:

# 3.1 <u>Description of data collection:</u>

Data are retrieved weekly.

## 3.2 <u>Description of derived parameters and processing techniques used:</u>

- 1) Soil temperature data were sampled every 5 seconds and their 30-minute averages were stored.
- 2) Soil moisture data were sampled every 5 minutes and their 30-minute averages were stored.
- 3) Soil moisture at 0.02 m was the average of output values from two sensors set horizontally in the soil at 2.5 cm depth.

# **4.0 QUALITY CONTROL PROCEDURES:**

At this stage of data processing, only apparently erroneous data were removed. Further quality control the data will be done later.

#### 5.0 GAP FILLING PROCEDURES:

<sup>&</sup>lt;sup>2</sup> The sensor was set obliquely in the soil.

<sup>&</sup>lt;sup>3</sup> Measurement was started at 2009/05/02 07:30 UTC.

At this stage of data processing, no gap filling procedure was applied. Gap filling will be done later.

### **6.0 DATA REMARKS:**

- 6.1 PI's assessment of the data:
- 6.1.1 <u>Instruments problems</u> None.

### 6.1.2 Quality issues

1) Soil moisture sensors were removed temporarily and set again at transplanting of rice. This caused abrupt changes in soil moisture data at 2009/05/02 07:30 (UTC).

```
6.2 Missing data periods:
from 2009/01/01 02:00 to 2009/01/04 02:30 (UTC) (Soil Temperature 0.02m)
from 2009/01/04 07:30 to 2009/01/04 10:30 (UTC) (Soil Temperature 0.02m)
from 2009/01/04 21:30 to 2009/01/05 06:30 (UTC) (Soil Temperature 0.02m)
from 2009/01/05 11:00 to 2009/01/05 13:30 (UTC) (Soil Temperature 0.02m)
from 2009/01/06 01:30 to 2009/01/06 06:30 (UTC) (Soil Temperature 0.02m)
from 2009/01/06 17:00 to 2009/01/07 04:30 (UTC) (Soil Temperature 0.02m)
from 2009/01/07 21:30 to 2009/01/08 04:00 (UTC) (Soil Temperature 0.02m)
2009/01/08 06:00 (UTC) (Soil Temperature 0.02m)
from 2009/01/09 12:30 to 2009/02/01 02:30 (UTC) (Soil Temperature 0.02m)
from 2009/02/01 06:30 to 2009/02/09 04:03 (UTC) (Soil Temperature 0.02m)
from 2009/02/01 05:00 to 2009/02/09 06:00 (UTC) (Soil Temperature 0.02m, Soil
     Moisture all depth)
2009/02/19 01:30 (UTC) (Soil Moisture 0.1m)
2009/02/19 02:30 (UTC) (Soil Moisture 0.1m)
2009/03/22 03:00 (UTC) (Soil Moisture 0.1m)
2009/03/22 04:30 (UTC) (Soil Moisture 0.1m)
2009/03/22 06:30 (UTC) (Soil Moisture 0.1m)
2009/03/22 08:30 (UTC) (Soil Moisture 0.1m)
2009/04/05 02:30 (UTC) (Soil Temperature all depth)
from 2009/04/20 06:00 to 2009/04/20 07:30 (UTC) (Soil Temperature 0.02m)
2009/04/21 5:00 (UTC) (Soil Temperature 0.02m)
from 2009/04/23 09:30 to 2009/04/23 14:30 (UTC) (Soil Temperature 0.02m)
2009/04/26 6:00 (UTC)(Soil Temperature all depth)
from 2009/04/26 06:30 to 2009/04/27 05:00 (UTC) (Soil Temperature all depth, Soil
     Moisture depth)
2009/04/27 5:30 (UTC) (Soil Temperature 0.02m, 0.1m)
from 2009/04/30 15:30 to 2009/05/01 08:00 (UTC) (Soil Temperature all depth)
2009/05/01 04:00 (UTC) (Soil Temperature 0.4m)
from 2009/05/01 05:30 to 2009/05/01 6:00 (UTC) (Soil Temperature 0.4m)
from 2009/05/01 08:30 to 2009/05/02 7:00 (UTC) (Soil Moisture all depth)
2009/05/04 14:30 (UTC) (Soil Temperature all depth)
from 2009/05/08 06:00 to 2009/05/08 13:00 (UTC) (Soil Temperature all depth)
from 2009/05/09 04:30 to 2009/05/12 07:30 (UTC) (Soil Temperature all depth)
```

from 2009/06/14 10:30 to 2009/06/15 01:30 (UTC) (Soil Temperature 0.02m) from 2009/06/15 05:30 to 2009/06/16 01:00 (UTC) (Soil Temperature 0.02m)

from 2009/06/16 13:00 to 2009/06/17 02:30 (UTC) (Soil Temperature 0.02m) from 2009/06/20 21:00 to 2009/06/23 02:00 (UTC) (Soil Temperature 0.02m) from 2009/06/23 23:00 to 2009/06/30 23:30 (UTC) (Soil Temperature 0.02m)

## **6.3 Data intercomparisons:**

# 7.0 REFERENCE REQUIREMENTS:

Original data were collected in the framework of Research Project for Global Warming Monitoring by NIAES. The project is funded by Ministry of Agriculture, Forestry and Fisheries, Ministry of Environment and NIAES.

### **8.0 REFERENCES**

- Saito, M, A. Miyata, H. Nagai, and T. Yamada, Seasonal variation of carbon dioxide exchange in rice paddy field in Japan. Agric. Forest Meteorol. 135, 93-109, 2005.
- Miyata, A., T. Iwata, H. Nagai, T. Yamada, H. Yoshikoshi, M. Mano, K. Ono, G. H. Han, Y. Harazono, E. Ohtaki, Md. A. Baten, S. Inohara, T. Takimoto, and M. Saito, Seasonal variation of carbon dioxide and methane fluxes at single cropping paddy fields in central and western Japan, Phyton, 45(4), 89-97, 2005.
- Saito, M., J. Asanuma, A. Miyata, Dual-scale transport of sensible heat and water vapor over a short canopy under unstable conditions. Water Resources Research, 43, W05413, doi:10.1029/2006WR005136, 2007.
- Han, G.H., H. Yoshikoshi, H. Nagai, T. Yamada, K. Ono, M. Mano, A. Miyata, Isotopic disequilibrium between carbon assimilated and respired in a rice paddy as influenced by methanogenesis from CO<sub>2</sub>. Journal of Geophysical Research, 112, G02016, doi:10.1029/2006JG000219, 2007.